

| Drug-Induced Death Rate | |
|--|--|
| SHIP website description | This indicator shows the drug-induced death rate per 100,000 population. Drug-induced deaths include all deaths for which illicit or prescription drugs are the underlying cause. In 2007, drug- induced deaths were more common than alcohol induced or firearm-related deaths in the United States. In 2010, there were 761 drug induced deaths in Maryland. |
| Source | Maryland MDH Vital Statistics Administration (VSA) |
| Numerator | Average number of deaths induced from drugs (Number of deaths with an ICD-10 code of D52.1, D59.0, D59.2, D61.1, D64.2, E06.4, E16.0, E23.1, E24.2, E27.3, E66.1, F11.0-F11.5, F11.7-F11.9, F12.0-F12.5, F12.7-F12.9, F13.0-F13.5, F13.7-F13.9, F14.0-F14.5, F14.7-F14.9, F15.0-F15.5, F15.7- F15.9, F16.0-F16.5, F16.7-F16.9, F17.0, F17.3-F17.5, F17.7-F17.9, F18.0-F18.5, F18.7-18.9, F19.0-F19.5, F19.7-F19.9, G21.1, G24.0, G25.1, G25.4, G25.6, G44.4, G62.0, G72.0, I95.2, J70.2-J70.4, K85.3, L10.5, L27.0-L27.1, M10.2, M32.0, M80.4, M81.4, M83.5, M87.1, R50.2, R78.1-R78.5, X40- X44, X60-X64, X85, Y10-14). Deaths by jurisdiction are by place of residence. |
| Denominator | Population (middle year) from Maryland MDH Vital Statistics Administration (VSA) |
| Threshold | 20 |
| Censorship (if below threshold) | Rate not reported if below threshold. Counts may be available upon request. |
| Calculation and metric | $(\text{Numerator}/\text{Denominator}) * 100,000 = 3\text{-Year Calculation}$. These data are age-adjusted to 2000 |
| Race/ethnicity categories (on SHIP chart) | All races/ ethnicities (aggregated) Black Non-Hispanic Hispanic White Non-Hispanic Asian/ Pacific Islander Non-Hispanic |
| Other | None |